NutriDyn

Brain Restore[,]

Nutritional Support for Cognitive Function*

Brain Restore Supplementation

NutriDyn's Brain Restore is a comprehensive formula that contains bioactive forms of key B vitamins, L-carnitine, choline, uridine, and serine to support cerebral and nervous system function.[•] Due to their role in supporting neurotransmitter production, these nutrients help support cognition, focus, and mood.[•]

Research cited herein suggests the nutrients contained in Brain Restore may support cerebral and nervous system function in a variety of ways.⁺ Moreover, these nutrients work in concert for proper DNA maintenance, energy production, amino acid metabolism, and a variety of other processes.⁺

- Supports cognitive function and healthy mood*
- Supports and maintains DNA⁺
- Supports blood and oxygen flow to the brain⁺
- Supports neurotransmitter production and neural tissue*
- Helps metabolize amino acids[•]

How Brain Restore Works





Brain Restore contains key B vitamins, including niacin, pyridoxine, folate, and vitamin B12 (cobalamin), which assist in the production of neurotransmitters.⁴ Longitudinal studies suggest deficiencies in several B vitamins accelerate brain tissue atrophy and cognitive decline.^{1,2}

For example, chronic deficiency of vitamin B12 can lead to serious, irreversible health consequences such as damage to the brain and nervous system and/or pernicious anemia (lack of red blood cell production).³ This is not surprising, given that vitamin B12 is needed for the body to convert homocysteine to methionine. Methionine is necessary for the formation of S-adenosylmethionine (SAMe), which is involved in the synthesis of catecholamines and various neurotransmitters.⁴

Recent research suggests that supplemental folic acid and vitamin B12 work synergistically to support cognition, energy production, and neural tissue.^{45,6} Pyridoxine is also important for cognition as it is necessary for the conversion of L-DOPA to dopamine, as well as the conversion of glutamate to GABA and the proper metabolism of L-tryptophan.⁴

Furthermore, niacin stimulates the production of a protein/growth factor in humans called brain-derived neurotrophic factor (BDNF). BDNF acts on neurons throughout the nervous system to encourage growth and replication, which is crucial for supporting long-term memory, learning, and overall cognition.⁴⁷

For more information, visit: www.nutridyn.com

How Brain Restore Works Continued

Brain Restore also contains acetyl-L-carnitine (ALCAR), a highly bioavailable form of L-carnitine. ALCAR can cross the blood-brain barrier, where it helps to manage oxidative damage, support mitochondrial function, and maintain normal neurotransmitter activity.⁴⁸

Rounding out the Brain Restore formula are Alpha-glycerolphosphorylcholine (Alpha-GPC), phosphatidyl serine, and uridine monophosphate (UMP). These phosphorylated compounds are readily absorbed by the body and support the production of neurotransmitters such as acetylcholine and dopamine.⁺ Studies suggest these nutrients work synergistically to support cognition and learning capacity.^{+ 9, 10}

Supplement Facts

Serving Size: 7 Capsules Servings Per Container: 30

	Amount Per Serving	%DV*
Niacin (as niacinamide)	100 mg NE	625%
Vitamin B6 (as pyridoxine HCI)	25 mg	1,470%
Folate (as calcium L-5-methyltetra	a- 1.7 mg DFE	425%
hydrofolate) (BioFolate®)		
Vitamin B12 (as methylcobalamin)) 2 mg	83,333%
Acetyl-L-Carnitine HCI	750 mg	**
Alpha-glycerolphosphorylcholine	(A-GPC) 600 mg	**
Uridine 5-Monophosphate	500 mg	**
Phosphatidyl Serine	150 mg	**

Other Ingredients: Hypromellose, sodium copper chlorophyllin, microcrystalline cellulose, tricalcium phosphate, vegetable magnesium stearate, silica.

BioFolate® is a federally registered trademark of MTC Industries, Inc.

Directions: Take seven capsules daily as a dietary supplement or as directed by your healthcare practitioner.

Caution: If you are pregnant, nursing, or taking medication, consult your healthcare practitioner before use. Keep out of reach of children.

References:

- Van Uffelen, J. G., Chinapaw, M. J., van Mechelen, W., & Hopman-Rock, M. (2008). Walking or vitamin B for cognition in older adults with mild cognitive impairment? A randomized controlled trial. *British journal of sports medicine*.
- Calvaresi, E., & Bryan, J. (2001). B Vitamins, Cognition, and Aging a Review. The Journals of Gerontology Series B: Psychological Sciences and Social Sciences, 56(6), P327-P339.
- HU, R. (2015). Vitamin B12 deficiency. European Journal of General Medicine, 12(3).
 Penninx, B. W., Guralnik, J. M., Ferrucci, L., Fried, L. P., Allen, R. H., & Stabler, S. P. (2000). Vitamin B12 deficiency and depression in physically disabled older women: epidemiologic evidence from the Women's Health and Aging Study. American Journal of Psychiatry, 157(5), 715-721.
- Coppen, A., & Bolander-Gouaille, C. (2005). Treatment of depression: time to consider folic acid and vitamin B12. *Journal of Psychopharmacology*, 19(1), 59-65.
- Eussen, S. J., de Groot, L. C., Joosten, L. W., Bloo, R. J., Clarke, R., Ueland, P. M., ... & van Staveren, W. A. (2006). Effect of oral vitamin B-12 with or without folic acid on cognitive function in older people with mild vitamin B-12 deficiency: a randomized, placebo-controlled trial. *The American journal of clinical nutrition*, 84(2), 361-370.
- Yamada K, Nabeshima T (Apr 2003). Brain-derived neurotrophic factor/TrkB signaling in memory processes. *Journal of Pharmacological Sciences*. 91(4): 267–70.
- Smeland, O. B., Meisingset, T. W., Borges, K., & Sonnewald, U. (2012). Chronic acetyl-L-carnitine alters brain energy metabolism and increases noradrenaline and serotonin content in healthy mice. *Neurochemistry international*. 61(1):100-10
- and serotonin content in healthy mice. Neurochemistry international, 61(1), 100-107.
 9. Holguin, S.; Martinez, J.; Chow, C.; Wurtman, R. (2008). "Dietary uridine enhances the improvement in learning and memory produced by administering DHA to gerbils". The FASEB Journal. 22(11): 3938–3946.
- Jorissen, B. L., Brouns, F. J. P. H., Van Boxtel, M. P. J., Ponds, R. W. H. M., Verhey, F. R. J., Jolles, J., & Riedel, W. J. (2001). The influence of soy-derived phosphatidylserine on cognition in age-associated memory impairment. *Nutritional neuroscience*, 4(2), 121-134.

• These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

For more information, visit: www.nutridyn.com